

COTANGENT BUNDLES WITH GENERAL NATURAL KÄHLER STRUCTURES

S. L. DRUȚĂ

We study the conditions under which an almost Hermitian structure (G, J) of general natural lift type on the cotangent bundle T^*M of a Riemannian manifold (M, g) is Kählerian. First, we obtain the algebraic conditions under which the manifold (T^*M, G, J) is almost Hermitian. Next we get the integrability conditions for the almost complex structure J , then the conditions under which the associated 2-form is closed. The manifold (T^*M, G, J) is Kählerian iff it is almost Kählerian and the almost complex structure J is integrable. It follows that the family of Kählerian structures of above type on T^*M depends on three essential parameters (one is a certain proportionality factor, the other two are parameters involved in the definition of J).

AMS 2000 Subject Classification: Primary 53C07, 53C15, 53C55.

Key words: cotangent bundle, Riemannian metric, general natural lift.

1. INTRODUCTION

The fundamental differences between the geometry of the cotangent bundle T^*M and that of the tangent bundle TM of a Riemannian manifold (M, g) , are due to the different construction of lifts to T^*M , which cannot be defined just like in the case of TM (see [14]).

The possibility to consider vertical, complete and horizontal lifts on T^*M leads to interesting geometric structures, studied in the last years (see [5], [8], [9], [10], [11], [12], [13], [14], [15], [16], [17], [18]). Due to the duality tangent bundle-cotangent bundle, some of these results are very much similar to previous results for the tangent bundles.

In the present paper we study the conditions under which a structure (G, J) of general natural lift type on the cotangent bundle T^*M of a Riemannian manifold M is a Kählerian structure. The similar problem for the tangent bundle was treated by Oproiu, in the paper [6] (see also [7] for the diagonal case). A part of the results from the papers [12], [15], [16], may be obtained as particular cases from those in the present paper.

First we consider a tensor field J of type $(1, 1)$ on T^*M which is a general natural lift of the Riemannian metric g . The condition for J to define an almost complex structure on T^*M leads to certain algebraic relations between the parameters involved in its definition. Four of the eight parameters involved in the definition of J may be obtained as (rational) functions of the other four parameters. The integrability condition for the almost complex structure J implies that the base manifold must have constant sectional curvature. Then some other relations fulfilled by the parameters and their derivatives are obtained, so that the essential parameters involved in the definition of the integrable almost complex structure are two.

In the next section we study the conditions under which a Riemannian metric G which is a general natural lift of g , is Hermitian with respect to J . We get that the structure (G, J) on T^*M is almost Hermitian if and only if the coefficients c_1, c_2, c_3 , involved in the definition of the metric G are

proportional to the coefficients a_1, a_2, a_3 , involved in definition of the almost complex structure J , with the proportionality factor λ , and the combinations $c_1 + 2td_1, c_2 + 2td_2, c_3 + 2td_3$ are proportional to $a_1 + 2tb_1, a_2 + 2tb_2, a_3 + 2tb_3$, the proportionality factor being chosen of the form $\lambda + 2t\mu$. The main result is that the structure (G, J) is Kählerian if and only if $\mu = \lambda'$ and the coefficients b_1, b_2, b_3 of the almost complex structure J may be expressed as certain rational functions of a_1, a_2, a_3 , and their derivatives. The condition for (G, J) to be Kählerian is obtained from the property for (G, J) to be almost Kählerian and the integrability property for J .

Some quite long computations have been done by using the Mathematica package RICCI for doing tensor calculations.

The manifolds, tensor fields and other geometric objects we consider in this paper are assumed to be differentiable of class C^∞ (i.e. smooth). We use the computations in local coordinates in a fixed local chart but many results may be expressed in an invariant form by using the vertical and horizontal lifts. The well known summation convention is used throughout this paper, the range of the indices h, i, j, k, l, m, r being always $\{1, \dots, n\}$.

2. PRELIMINARY RESULTS

Let (M, g) be a smooth n -dimensional Riemannian manifold and denote its cotangent bundle by $\pi : T^*M \longrightarrow M$. Recall that there is a structure of a $2n$ -dimensional smooth manifold on T^*M , induced from the structure of smooth n -dimensional manifold of M . From every local chart $(U, \varphi) = (U, x^1, \dots, x^n)$ on M , it is induced a local chart $(\pi^{-1}(U), \Phi) = (\pi^{-1}(U), q^1, \dots, q^n, p_1, \dots, p_n)$, on T^*M , as follows. For a cotangent vector $p \in \pi^{-1}(U) \subset T^*M$, the first n local coordinates $q^1, \dots, q^n$ are the local coordinates of its base point $x = \pi(p)$ in the local chart (U, φ) (in fact we have $q^i = \pi^*x^i = x^i \circ \pi, i = 1, \dots, n$). The last n local coordinates $p_1, \dots, p_n$ of $p \in \pi^{-1}(U)$ are the vector space coordinates of p with respect to the natural basis $(dx_{\pi(p)}^1, \dots, dx_{\pi(p)}^n)$, defined by the local chart (U, φ) , i.e. $p = p_i dx_{\pi(p)}^i$.

An M -tensor field of type (r, s) on T^*M is defined by sets of n^{r+s} components (functions depending on q^i and p_i), with r upper indices and s lower indices, assigned to induced local charts $(\pi^{-1}(U), \Phi)$ on T^*M , such that the local coordinate change rule is that of the local coordinate components of a tensor field of type (r, s) on the base manifold M (see [4] for further details in the case of the tangent bundle). An usual tensor field of type (r, s) on M may be thought as an M -tensor field of type (r, s) on T^*M . If the considered tensor field on M is covariant only, the corresponding M -tensor field on T^*M may be identified with the induced (pullback by π) tensor field on T^*M .

Some useful M -tensor fields on T^*M may be obtained as follows. Let $v, w : [0, \infty) \longrightarrow \mathbf{R}$ be smooth functions and let $\|p\|^2 = g_{\pi(p)}^{-1}(p, p)$ be the square of the norm of the cotangent vector $p \in \pi^{-1}(U)$ (g^{-1} is the tensor field of type $(2, 0)$ having the components $(g^{kl}(x))$, which are the entries of the inverse of the matrix $(g_{ij}(x))$, defined by the components of g in the local chart (U, φ)). The components $vg_{ij}(\pi(p)), p_i, w(\|p\|^2)p_i p_j$ define respective M -tensor fields of types $(0, 2), (0, 1), (0, 2)$ on T^*M . Similarly, the components $vg^{kl}(\pi(p)), g^{0i} = p_h g^{hi}, w(\|p\|^2)g^{0k}g^{0l}$ define respective M -tensor fields of type $(2, 0), (1, 0), (2, 0)$ on T^*M . Of course, all the above components are considered in the induced local chart $(\pi^{-1}(U), \Phi)$.

The Levi Civita connection $\dot{\nabla}$ of g defines a direct sum decomposition

$$(2.1) \quad TT^*M = VT^*M \oplus HT^*M.$$

of the tangent bundle to T^*M into vertical distributions $VT^*M = \text{Ker } \pi_*$ and the horizontal distribution HT^*M .

If $(\pi^{-1}(U), \Phi) = (\pi^{-1}(U), q^1, \dots, q^n, p_1, \dots, p_n)$ is a local chart on T^*M , induced from the local chart $(U, \varphi) = (U, x^1, \dots, x^n)$, the local vector fields $\frac{\partial}{\partial p_1}, \dots, \frac{\partial}{\partial p_n}$ on $\pi^{-1}(U)$ define a local frame for VT^*M over $\pi^{-1}(U)$ and the local vector fields $\frac{\delta}{\delta q^1}, \dots, \frac{\delta}{\delta q^n}$ define a local frame for HT^*M over $\pi^{-1}(U)$, where

$$\frac{\delta}{\delta q^i} = \frac{\partial}{\partial q^i} + \Gamma_{ih}^0 \frac{\partial}{\partial p_h}, \quad \Gamma_{ih}^0 = p_k \Gamma_{ih}^k$$

and $\Gamma_{ih}^k(\pi(p))$ are the Christoffel symbols of g .

The set of vector fields $\{\frac{\partial}{\partial p_1}, \dots, \frac{\partial}{\partial p_n}, \frac{\delta}{\delta q^1}, \dots, \frac{\delta}{\delta q^n}\}$ defines a local frame on T^*M , adapted to the direct sum decomposition (2.1).

We consider

$$(2.2) \quad t = \frac{1}{2} \|p\|^2 = \frac{1}{2} g_{\pi(p)}^{-1}(p, p) = \frac{1}{2} g^{ik}(x) p_i p_k, \quad p \in \pi^{-1}(U),$$

the energy density defined by g in the cotangent vector p . We have $t \in [0, \infty)$ for all $p \in T^*M$.

Roughly speaking, the natural lifts have coefficients as functions of the density energy only ([1], [2], [3]).

From now on we shall work in a fixed local chart (U, φ) on M and in the induced local chart $(\pi^{-1}(U), \Phi)$ on T^*M .

We may easily obtain the following

LEMMA 2.1. *If $n > 1$ and u, v are smooth functions on T^*M such that either*

$$ug_{ij} + vp_i p_j = 0, \quad ug^{ij} + vg^{0i} g^{0j} = 0, \quad \text{or} \quad u\delta_j^i + vg^{0i} p_j = 0$$

*on the domain of any induced local chart on T^*M , then $u = 0$, $v = 0$.*

3. THE INTEGRABILITY OF THE ALMOST COMPLEX STRUCTURES

If X is a vector field $X \in \mathcal{X}(M)$, then g_X is the 1-form on M defined by the relation $g_X(Y) = g(X, Y)$, $\forall Y \in \mathcal{X}(M)$. Using the local chart of M , $(U, x^1, \dots, x^n)$, we may write $X = X^i \frac{\partial}{\partial x^i}$. Then $g_X = g_{ij} X^j dx^i$, $g_{\frac{\partial}{\partial x^i}} = g_{ij} dx^j$.

Let us consider a 1-form $\theta \in \Lambda^1(M)$. Then $\theta^\sharp = g_\theta^{-1}$ is a vector field on M defined by the musical isomorphism $g(\theta^\sharp, Y) = \theta(Y)$, $\forall Y \in \mathcal{X}(M)$. If $\theta = \theta_i dx^i$, then $\theta^\sharp = g^{ij} \theta_j \frac{\partial}{\partial x^i}$, the expressions being on the local chart $(U, x^1, \dots, x^n)$ of M .

For $p \in T^*M$ we consider the vector $p^\sharp$ tangent to M in $\pi(p)$.

The Liouville vector field on T^*M is p^V , expressed by $p_p^V = p_i \frac{\partial}{\partial p_i}$ in every point p of the induced local chart $(\pi^{-1}(U), \Phi)$ on T^*M .

The similar horizontal vector field on T^*M is $(p^\sharp)^H$, with the expression $(p_p^\sharp)^H = g^{0i} \frac{\delta}{\delta q^i}$ in every point p of the induced local chart $(\pi^{-1}(U), \Phi)$ on T^*M (recall that $g^{0i} = p_h g^{hi}$).

Consider the real valued smooth functions $a_1, a_2, a_3, a_4, b_1, b_2, b_3, b_4$ defined on $[0, \infty) \subset \mathbf{R}$. We define a *general natural tensor of type (1, 1)* on T^*M by its action on the horizontal and vertical lifts on T^*M

$$\begin{cases} JX_p^H = a_1(t)(g_X)_p^V + b_1(t)p(X)p_p^V + a_4(t)X_p^H + b_4(t)p(X)(p^\sharp)_p^H, \\ J\theta_p^V = a_3(t)\theta_p^V + b_3(t)g_{\pi(p)}^{-1}(p, \theta)p_p^V - a_2(t)(\theta^\sharp)_p^H - b_2(t)g_{\pi(p)}^{-1}(p, \theta)(p^\sharp)_p^H, \end{cases}$$

in every point p of the induced local card $(\pi^{-1}(U), \Phi)$ on T^*M , $\forall X \in \mathcal{X}(M), \forall \theta \in \Lambda^1(M)$.

With respect to the local adapted frame $\{\frac{\delta}{\delta q^i}, \frac{\partial}{\partial p_j}\}_{i,j=1,\dots,n}$, the expressions of J become

$$(3.1) \quad \begin{cases} J\frac{\delta}{\delta q^i} = a_1(t)g_{ij}\frac{\partial}{\partial p_j} + b_1(t)p_iC + a_4(t)\frac{\delta}{\delta q^i} + b_4(t)p_i\tilde{C}, \\ J\frac{\partial}{\partial p_i} = a_3(t)\frac{\partial}{\partial p_i} + b_3(t)g^{0i}C - a_2(t)g^{ij}\frac{\delta}{\delta q^j} - b_2(t)g^{0i}\tilde{C}, \end{cases}$$

where we have denoted by $C = p^V$ the Liouville vector-field on T^*M and by $\tilde{C} = (p^\sharp)^H$, the corresponding horizontal vector field on T^*M .

We may write also

$$(3.2) \quad \begin{cases} J\frac{\delta}{\delta q^i} = J_{ij}^{(1)}\frac{\partial}{\partial p_j} + J4_i^j\frac{\delta}{\delta q^j}, \\ J\frac{\partial}{\partial p_i} = J3_j^i\frac{\partial}{\partial p_j} - J_{(2)}^{ij}\frac{\delta}{\delta q^j}, \end{cases}$$

where

$$J_{ij}^{(1)} = a_1(t)g_{ij} + b_1(t)p_ip_j, \quad J4_i^j = a_4(t)\delta_i^j + b_4(t)g^{0j}p_i,$$

$$J3_j^i = a_3(t)\delta_j^i + b_3(t)g^{0i}p_j, \quad J_{(2)}^{ij} = a_2(t)g^{ij} + b_2(t)g^{0i}g^{0j}.$$

THEOREM 3.1. *A natural tensor field J of type (1, 1) on T^*M given by (3.1) or (3.2) defines an almost complex structure on T^*M , if and only if $a_4 = -a_3, b_4 = -b_3$ and the coefficients a_1, a_2, a_3, b_1, b_2 and b_3 are related by*

$$(3.3) \quad a_1a_2 = 1 + a_3^2, \quad (a_1 + 2tb_1)(a_2 + 2tb_2) = 1 + (a_3 + 2tb_3)^2.$$

Remark. From the conditions (3.3) we get that the coefficients $a_1, a_2, a_1 + 2tb_1, a_2 + 2tb_2$ have the same sign and cannot vanish. We assume that $a_1 > 0, a_2 > 0, a_1 + 2tb_1 > 0, a_2 + 2tb_2 > 0$ for all $t \geq 0$.

Remark. The relations (3.3) allow us to express two of the coefficients $a_1, a_2, a_3, b_1, b_2, b_3$, as functions of the other four; e.g. we have:

$$(3.4) \quad a_2 = \frac{1 + a_3^2}{a_1}, \quad b_2 = \frac{2a_3b_3 - a_2b_1 + 2tb_3^2}{a_1 + 2tb_1}.$$

The integrability condition for the above almost complex structure J on a manifold M is characterized by the vanishing of its Nijenhuis tensor field N_J , defined by

$$N_J(X, Y) = [JX, JY] - J[JX, Y] - J[X, JY] - [X, Y],$$

for all vector fields X and Y on M .

THEOREM 3.2. *Let (M, g) be an $n(> 2)$ -dimensional connected Riemannian manifold. The almost complex structure J defined by (3.1) on T^*M is integrable if and only if (M, g) has constant sectional curvature c and the coefficients b_1, b_2, b_3 are given by:*

$$(3.5) \quad \begin{cases} b_1 = \frac{2c^2ta_2^2 + 2cta_1a_2' + a_1a_1' - c + 3ca_3^2}{a_1 - 2ta_1' - 2cta_2 - 4ct^2a_2'}, \\ b_2 = \frac{2ta_3'^2 - 2ta_1'a_2' + ca_2^2 + 2cta_2a_2' + a_1a_2'}{a_1 - 2ta_1' - 2cta_2 - 4ct^2a_2'}, \\ b_3 = \frac{a_1a_3' + 2ca_2a_3 + 4cta_2'a_3 - 2cta_2a_3'}{a_1 - 2ta_1' - 2cta_2 - 4ct^2a_2'}. \end{cases}$$

Proof. From the condition $N_J(\frac{\partial}{\partial p_i}, \frac{\partial}{\partial p_j}) = 0$ we obtain that the horizontal component of this Nijenhuis bracket vanishes if and only if

$$(3.6) \quad a_2' = \frac{a_2a_3' + 2a_3b_2 - a_2b_3}{2(a_3 + tb_3)}$$

and the vertical component vanishes if and only if

$$(3.7) \quad (a_1a_2' - a_1b_2 + 2a_3'b_3t)(\delta_j^h p_i - \delta_i^h p_j) - a_2^2 g^{0k} R_{kij}^h - a_2 b_2 g^{0k} g^{0l} (p_j R_{kil}^h - p_i R_{kjl}^h) = 0.$$

Taking into account that the curvature of the base manifold does not depend on p , we differentiate with respect to p_k in (3.7). Considering the value of this derivative in $p = 0$, we get

$$(3.8) \quad R_{kij}^h = c(\delta_i^h g_{kj} - \delta_j^h g_{ki}),$$

where

$$c = \frac{a_1(0)}{a_2^2(0)}(b_2(0) - a_2'(0)),$$

which is a function depending on $q^1, \dots, q^n$ only. According to the Schur's theorem, c must be a constant when $n > 2$ and M is connected.

Using the condition of constant sectional curvature for the base manifold, we obtain from $N_J(\frac{\delta}{\delta q^i}, \frac{\delta}{\delta q^j}) = 0$, the following expressions for a_1' and a_3'

$$(3.9) \quad \begin{cases} a_1' = \frac{a_1b_1 + c(1 - 3a_2^2 - 4ta_3b_3)}{a_1 + 2tb_1}, \\ a_3' = \frac{a_1b_3 - 2ca_2(a_3 + tb_3)}{a_1 + 2tb_1}. \end{cases}$$

If we replace in (3.6) the expression of a_3' and the relations (3.4), we may write a_2' in the form

$$(3.10) \quad a_2' = \frac{2a_3b_3 - a_2b_1 - ca_2^2}{a_1 + 2tb_1}.$$

The values of a_1' , a_3' in (3.9) and a_2' in (3.10) fulfill the vanishing condition for the vertical component of the Nijenhuis bracket $N_J(\frac{\partial}{\partial p_i}, \frac{\delta}{\delta q^j})$. The same expressions fulfill also the relation

$$(3.11) \quad a_1a_2' + a_1'a_2 = 2a_3a_3',$$

obtained by differentiating the first of the relation (3.3) with respect to t .

We may solve the system given by (3.6) and (3.9), with respect to b_1, b_2, b_3 . Taking (3.11) into account, we get the relations (3.5) from the theorem, which fulfill identically the expression of b_2 from (3.4).

Remark. In the diagonal case, where $a_3 = 0$ it follows $b_3 = 0$ too, and we have:

$$a_2 = \frac{1}{a_1}, \quad b_1 = \frac{a_1 a_1' - c}{a_1 - 2ta_1'}, \quad b_2 = \frac{c - a_1 a_1'}{a_1(a_1^2 - 2ct)}.$$

Thus we found some results from [13] and [16], papers which have treated only the diagonal case.

4. KÄHLER STRUCTURES OF GENERAL NATURAL LIFT TYPE ON THE COTANGENT BUNDLE

In this section, we introduce a *Riemannian metric* G of general natural lift type on the cotangent bundle T^*M , defined by

$$(4.1) \quad \begin{cases} G_p(X^H, Y^H) = c_1(t)g_{\pi(p)}(X, Y) + d_1(t)p(X)p(Y), \\ G_p(\theta^V, \omega^V) = c_2(t)g_{\pi(p)}^{-1}(\theta, \omega) + d_2(t)g_{\pi(p)}^{-1}(p, \theta)g_{\pi(p)}^{-1}(p, \omega), \\ G_p(X^H, \theta^V) = G_p(\theta^V, X^H) = c_3(t)\theta(X) + d_3(t)p(X)g_{\pi(p)}^{-1}(p, \theta), \end{cases}$$

$\forall X, Y \in \mathcal{X}(M), \forall \theta, \omega \in \Lambda^1(M), \forall p \in T^*M$.

The expressions (4.1) in local coordinates become

$$(4.2) \quad \begin{cases} G(\frac{\delta}{\delta q^i}, \frac{\delta}{\delta q^j}) = c_1(t)g_{ij} + d_1(t)p_i p_j = G_{ij}^{(1)}, \\ G(\frac{\partial}{\partial p_i}, \frac{\partial}{\partial p_j}) = c_2(t)g^{ij} + d_2(t)g^{0i}g^{0j} = G_{(2)}^{ij}, \\ G(\frac{\partial}{\partial p_i}, \frac{\delta}{\delta q^j}) = G(\frac{\delta}{\delta q^i}, \frac{\partial}{\partial p_j}) = c_3(t)\delta_i^j + d_3(t)p_i g^{0j} = G3_i^j, \end{cases}$$

where $c_1, c_2, c_3, d_1, d_2, d_3$ are six smooth functions of the density energy on T^*M . The conditions for G to be positive definite are assured if

$$(4.3) \quad \begin{aligned} c_1 + 2td_1 &> 0, \quad c_2 + 2td_2 > 0, \\ (c_1 + 2td_1)(c_2 + 2td_2) - (c_3 + 2td_3)^2 &> 0. \end{aligned}$$

The metric G is almost Hermitian with respect to the general almost complex structure J , if

$$G(JX, JY) = G(X, Y),$$

for all vector fields X, Y on T^*M .

We prove the following result

THEOREM 4.1. *The family of natural Riemannian metrics G on T^*M such that (T^*M, G, J) is an almost Hermitian manifold, is given by (4.2), provided that the coefficients c_1, c_2, c_3, d_1, d_2 , and d_3 are related to the coefficients a_1, a_2, a_3, b_1, b_2 , and b_3 by the following proportionality relations*

$$(4.4) \quad \frac{c_1}{a_1} = \frac{c_2}{a_2} = \frac{c_3}{a_3} = \lambda$$

$$(4.5) \quad \frac{c_1 + 2td_1}{a_1 + 2tb_1} = \frac{c_2 + 2td_2}{a_2 + 2tb_2} = \frac{c_3 + 2td_3}{a_3 + 2tb_3} = \lambda + 2t\mu,$$

where the proportionality coefficients $\lambda > 0$ and $\lambda + 2t\mu > 0$ are functions depending on t .

Proof. We use the local adapted frame $\{\frac{\delta}{\delta q^i}, \frac{\partial}{\partial p_j}\}_{i,j=1,\dots,n}$. The metric G is almost hermitian, if and only if the following conditions are fulfilled

$$(4.6) \quad \begin{cases} G(J\frac{\delta}{\delta q^i}, J\frac{\delta}{\delta q^j}) = G(\frac{\delta}{\delta q^i}, \frac{\delta}{\delta q^j}), \\ G(J\frac{\partial}{\partial p_i}, J\frac{\partial}{\partial p_j}) = G(\frac{\partial}{\partial p_i}, \frac{\partial}{\partial p_j}), \\ G(J\frac{\partial}{\partial p_i}, J\frac{\delta}{\delta q^j}) = G(\frac{\partial}{\partial p_i}, \frac{\delta}{\delta q^j}). \end{cases}$$

By using Lemma 2.1 we obtain that the coefficients of g_{ij} , g^{ij} , δ_j^i in the conditions (4.6) must vanish. It follows that the parameters c_1 , c_2 , c_3 , from the definition of the metric G , satisfy the homogeneous linear system of the form

$$(4.7) \quad \begin{cases} (a_3^2 - 1)c_1 + a_1^2 c_2 - 2a_1 a_3 c_3 = 0, \\ a_2^2 c_1 + (a_3^2 - 1)c_2 - 2a_2 a_3 c_3 = 0, \\ a_2 a_3 c_1 + a_1 a_3 c_2 - 2a_1 a_2 c_3 = 0. \end{cases}$$

The nontrivial solutions of (4.7) are given by the expression (4.4).

From the vanishing condition for the coefficients of $p_i p_j$, $g^{0i} g^{0j}$, $g^{0i} p_j$ in (4.6), we obtain a much more complicated system, fulfilled by d_1 , d_2 , d_3 . In order to get a certain similitude with the above system (4.7), fulfilled by c_1 , c_2 , c_3 , we multiply the new equations by $2t$ and substract the equations of the system (4.7), respectively. The new system may be written in a form in which the new unknowns are $c_1 + 2td_1$, $c_2 + 2td_2$, $c_3 + 2td_3$:

$$\begin{cases} [(a_3 + 2tb_3)^2 - 1](c_1 + 2td_1) + (a_1 + 2tb_1)^2(c_2 + 2td_2) \\ - 2(a_1 + 2tb_1)(a_3 + 2tb_3)(c_3 + 2td_3) = 0, \\ (a_2 + 2tb_2)^2(c_1 + 2td_1) + [(a_3 + 2tb_3)^2 - 1](c_2 + 2td_2) \\ - 2(a_2 + 2tb_2)(a_3 + 2tb_3)(c_3 + 2td_3) = 0, \\ (a_2 + 2tb_2)(a_3 + 2tb_3)(c_1 + 2td_1) + (a_1 + 2tb_1)(a_3 + 2tb_3)(c_2 + 2td_2) \\ - 2(a_1 + 2tb_1)(a_2 + 2tb_2)(c_3 + 2td_3) = 0. \end{cases}$$

Then the nonzero solutions are given by the relation (4.5).

The conditions (4.3) are fullfield, due to the properties (3.4) of the coefficients of the almost complex structure J .

The explicit expressions of the coefficients d_1, d_2, d_3 , obtained from (4.5), are

$$(4.8) \quad \begin{cases} d_1 = \lambda b_1 + \mu(a_1 + 2tb_1), \\ d_2 = \lambda b_2 + \mu(a_2 + 2tb_2), \\ d_3 = \lambda b_3 + \mu(a_3 + 2tb_3). \end{cases}$$

Remark. In the case when $a_3 = 0$, it follows that $c_3 = d_3 = 0$ and we obtain the almost Hermitian structure considered in [13] and [16]. Moreover, if $\lambda = 1$ and $\mu = 0$, we obtain the almost Kählerian structure considered in the mentioned papers.

Considering the two-form Ω defined by the almost Hermitian structure (G, J) on T^*M

$$\Omega(X, Y) = G(X, JY),$$

for all vector fields X, Y on T^*M , we obtain the following result:

PROPOSITION 4.2. *The expression of the 2-form Ω in the local adapted frame $\{\frac{\delta}{\delta q^i}, \frac{\partial}{\partial p_j}\}_{i,j=1,\dots,n}$ on T^*M , is given by*

$$\Omega\left(\frac{\partial}{\partial p_i}, \frac{\partial}{\partial p_j}\right) = 0, \quad \Omega\left(\frac{\delta}{\delta q^i}, \frac{\delta}{\delta q^j}\right) = 0, \quad \Omega\left(\frac{\partial}{\partial p_i}, \frac{\delta}{\delta q^j}\right) = \lambda \delta_j^i + \mu g^{0i} p_j$$

or, equivalently

$$(4.9) \quad \Omega = (\lambda \delta_j^i + \mu g^{0i} p_j) Dp_i \wedge dq^j,$$

where $Dp_i = dp_i - \Gamma_{ih}^0 dq^h$ is the absolute differential of p_i .

Next, by calculating the exterior differential of Ω , we obtain:

THEOREM 4.3. *The almost Hermitian structure (T^*M, G, J) is almost Kählerian if and only if*

$$\mu = \lambda'.$$

Proof. The differential of Ω is

$$d\Omega = (d\lambda \delta_j^i + d\mu g^{0i} p_j + \mu dg^{0i} p_j + \mu g^{0i} dp_j) \wedge Dp_i \wedge dq^j + (\lambda \delta_j^i + \mu g^{0i} p_j) dDp_i \wedge dq^j.$$

We first obtain the expressions of $d\lambda$, $d\mu$, dg^{0i} and dDp_i :

$$d\lambda = \lambda' g^{0h} Dp_h, \quad d\mu = \mu' g^{0h} Dp_h, \quad dg^{0i} = g^{hi} Dp_h - \Gamma_{j0}^i dq^j,$$

$$dDp_i = \frac{1}{2} R_{ijh}^0 dq^h \wedge dq^j - \Gamma_{ij}^h Dp_h \wedge dq^j.$$

By substituting these relations in $d\Omega$, using the properties of the external product, the symmetry of g^{ij} , Γ_{ij}^h and the Bianchi identities, we get

$$d\Omega = \frac{1}{2} (\lambda' - \mu) p_k (g^{kh} \delta_j^i - g^{ki} \delta_j^h) Dp_h \wedge Dp_i \wedge dq^j.$$

Hence $d\Omega = 0$ if and only if $\mu = \lambda'$.

Remark. Thus the family of general natural almost Kählerian structures on T^*M depends on five essential coefficients a_1 , a_3 , b_1 , b_3 , λ , which must satisfy the supplementary conditions $a_1 > 0$, $a_1 + 2tb_1 > 0$, $\lambda > 0$, $\lambda + 2t\mu > 0$.

Combining the results from the theorems 4.1, 3.2 and 4.3 we may state

THEOREM 4.4. *An almost Hermitian structure (G, J) of general natural lift type on T^*M is Kählerian if and only if the almost complex structure J is integrable (see Theorem 3.2) and $\mu = \lambda'$.*

Remark. The family of general natural Kählerian structures on T^*M depends on three essential coefficients a_1 , a_3 , λ , which must satisfy the supplementary conditions $a_1 > 0$, $a_1 + 2tb_1 > 0$, $\lambda > 0$, $\lambda + 2t\lambda' > 0$, where b_1 is given by (3.5).

Examples of such structures may be found in [12], [13], [16].

Acknowledgements. The author expresses her gratitude to her PhD adviser, professor V.Oproiu, for several discussions on this argument, several hints and encouragements.

The research was partially supported by the Grant, TD 158/2007, CNCSIS, Ministerul Educației și Cercetării, România.

REFERENCES

- [1] I. Kolář, P. Michor, J. Slovák, *Natural Operations in Differential Geometry*. Springer Verlag, Berlin, 1993, vi, 434 pp.
- [2] O. Kowalski, M. Sekizawa, *Natural Transformations of Riemannian Metrics on Manifolds to Metrics on Tangent Bundles-a Classification*. Bull. Tokyo Gakugei Univ. (4), **40** (1988), 1-29.
- [3] D. Krupka, J. Janyška, *Lectures on Differential Invariants*. Folia Fac. Sci. Nat. Univ. Purkiniane Brunensis, 1990.
- [4] K.P. Mok, E.M. Patterson, Y.C. Wong *Structure of Symmetric Tensors of Type (0,2) and Tensors of Type (1,1) on the Tangent Bundle*. Trans. Amer. Math. Soc., **234** (1977), 253-278.
- [5] M.I. Munteanu, *CR-structures on the Unit Cotangent Bundle*, An. Șt. Univ. "Al. I. Cuza" Iași, Math., **44** (1998), s.I, f.1, 125-136.
- [6] V. Oproiu, *A Generalization of Natural Almost Hermitian Structures on the Tangent Bundles*. Math. J. Toyama Univ. **22** (1999) 1-14.
- [7] V. Oproiu, *Some New Geometric Structures on the Tangent Bundles*. Publ. Math. Debrecen, **55/3-4** (1999), 261-281.
- [8] V. Oproiu, N. Papaghiuc, *A Pseudo-Riemannian Structure on the Cotangent Bundle*. An. Șt. Univ. "Al. I. Cuza" Iași, **36** (1990), 265-276.
- [9] V. Oproiu, N. Papaghiuc, *Another Pseudo-Riemannian Structure on the Cotangent Bundle*. Blt. Instit. Politehn. Iași **37** (41), Fasc. 1-4 (1991), Sect. I, 27-33.
- [10] V. Oproiu, N. Papaghiuc, *Locally Symmetric Cotangent Bundles*. Matematicki Vesnik, **42** (1990), 221-232.
- [11] V. Oproiu, N. Papaghiuc, G. Mitric, *Some Classes of Parahermitian Structures on cotangent bundles*. An. Șt. Univ. "Al. I. Cuza" Iași **43**, s.I.a, Matematică (1997), f1, 7-22.
- [12] V. Oproiu, D.D. Poroșniuc, *A class of Kaehler Einstein Structures on the Cotangent Bundle of a Space Form*. Publ. Math. Debrecen **66** (2005), 457-478.
- [13] V. Oproiu, D.D. Poroșniuc, *A Kähler Einstein Structure on the Cotangent Bundle of a Riemannian Manifold*. An. Științ. Univ. Al. I. Cuza, Iași **49**, s.I, Mathematics (2003), f.2, 399-414.
- [14] D.D. Poroșniuc, *A Class of Locally Symmetric Kähler Einstein Structures on the Nonzero Cotangent Bundle of a Space Form*. Balkan Journal of Geometry and Its Applications, **9** (2004), no.2, 68-81.
- [15] D.D. Poroșniuc, *A Class of Kähler Einstein Structures on the Nonzero Cotangent Bundle of a Space Form*. Rev. Roumaine Math. Pures Appl. **50** (2005), no. 2, 237-252.
- [16] D.D. Poroșniuc, *A Kähler Einstein Structure on the Nonzero Cotangent Bundle of a Space Form*. Ital. J. Pure Appl. Math. No. **18** (2005), 223-234.
- [17] D.D. Poroșniuc, *A Locally Symmetric Kähler Einstein Structure on the Cotangent Bundle of a Space Form*. Balkan Journal of Geometry and Its Applications, **9** (2004), no.1, 94-103.
- [18] D.D. Poroșniuc, *A Locally Symmetric Kähler Einstein Structure on a Tube in the Nonzero Cotangent Bundle of a Space Form*. An. Științ. Univ. Al. I. Cuza Iași. Mat. (N.S.) **50** (2004), no. 2, 315-326 (2005).
- [19] K. Yano, S. Ishihara, *Tangent and Cotangent Bundles*. M. Dekker Inc., New York, 1973.

"Al.I. Cuza" University,
 Faculty of Mathematics,
 Bd. Carol I, Nr. 11,
 Iași, 700 506, ROMÂNIA
 simonadruta@yahoo.com